

Armenia - The probabilities of a hip fracture in men

The following tables give the 10-year probability (%) of a hip fracture according to the T-score for femoral neck BMD , the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Armenia. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Note that the BMI is set at 24 kg/m².

Age = 50 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	14	7.9	4.2	2.2	1.2	0.6	0.3	0.2	0.1	0.1	0.0
1	22 (15-28)	12 (8.3-16)	6.7 (4.4-8.8)	3.6 (2.4-4.7)	1.9 (1.2-2.5)	1.0 (0.7-1.3)	0.5 (0.3-0.7)	0.3 (0.2-0.4)	0.1 (0.1-0.2)	0.1 (0.1-0.1)	0.0 (0.0-0.1)
2	31 (21-43)	18 (12-26)	10 (6.3-15)	5.6 (3.3-8.3)	3.0 (1.8-4.5)	1.6 (0.9-2.4)	0.8 (0.5-1.3)	0.4 (0.3-0.7)	0.2 (0.1-0.4)	0.1 (0.1-0.2)	0.1 (0.0-0.1)
3	43 (29-57)	27 (17-38)	15 (9.4-23)	8.5 (5.1-13)	4.6 (2.7-7.2)	2.5 (1.4-3.9)	1.3 (0.8-2.1)	0.7 (0.4-1.1)	0.4 (0.2-0.6)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
4	56 (43-69)	37 (26-51)	23 (15-33)	13 (8.2-19)	7.0 (4.4-11)	3.8 (2.4-5.9)	2.0 (1.2-3.1)	1.1 (0.7-1.7)	0.6 (0.4-0.9)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
5	68 (59-77)	50 (40-61)	32 (24-42)	19 (14-26)	11 (7.7-15)	5.8 (4.1-8.2)	3.1 (2.2-4.4)	1.6 (1.2-2.4)	0.9 (0.6-1.3)	0.5 (0.3-0.7)	0.3 (0.2-0.4)
6	78	63	44	27	16	8.6	4.7	2.5	1.3	0.7	0.4

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Armenia.

Age = 55 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	14	7.8	4.3	2.4	1.3	0.7	0.4	0.2	0.1	0.1	0.0
1	20 (14-25)	12 (8.1-15)	6.6 (4.6-8.3)	3.7 (2.5-4.7)	2.1 (1.4-2.6)	1.1 (0.8-1.4)	0.6 (0.4-0.8)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)	0.1 (0.0-0.1)
2	28 (20-38)	17 (11-24)	10 (6.4-14)	5.7 (3.6-8.1)	3.2 (2.0-4.5)	1.7 (1.1-2.5)	1.0 (0.6-1.4)	0.5 (0.3-0.8)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)
3	39 (28-50)	25 (17-34)	15 (9.6-21)	8.5 (5.4-12)	4.8 (3.0-7.0)	2.7 (1.7-4.0)	1.5 (0.9-2.2)	0.8 (0.5-1.2)	0.5 (0.3-0.7)	0.3 (0.2-0.4)	0.1 (0.1-0.2)
4	50 (39-62)	34 (25-45)	21 (15-29)	12 (8.5-18)	7.1 (4.8-11)	4.0 (2.7-6.0)	2.2 (1.5-3.3)	1.2 (0.8-1.9)	0.7 (0.5-1.1)	0.4 (0.3-0.6)	0.2 (0.2-0.3)
5	62 (54-71)	45 (37-55)	29 (24-38)	18 (14-24)	11 (8.1-15)	6.0 (4.6-8.4)	3.3 (2.5-4.7)	1.9 (1.4-2.6)	1.1 (0.8-1.5)	0.6 (0.5-0.9)	0.3 (0.3-0.5)
6	72	57	40	25	15	8.8	4.9	2.8	1.6	0.9	0.5

Age = 60 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	11	6.5	3.9	2.3	1.3	0.8	0.5	0.3	0.2	0.1	0.1
1	16 (11-19)	9.6 (6.9-11)	5.8 (4.1-6.9)	3.4 (2.4-4.1)	2.0 (1.4-2.4)	1.2 (0.8-1.4)	0.7 (0.5-0.8)	0.4 (0.3-0.5)	0.2 (0.2-0.3)	0.1 (0.1-0.2)	0.1 (0.1-0.1)
2	22 (16-28)	14 (9.6-18)	8.5 (5.7-11)	5.1 (3.4-6.9)	3.0 (2.0-4.1)	1.8 (1.2-2.4)	1.0 (0.7-1.4)	0.6 (0.4-0.8)	0.4 (0.2-0.5)	0.2 (0.2-0.3)	0.1 (0.1-0.2)
3	30 (22-38)	19 (14-26)	12 (8.6-17)	7.4 (5.1-10)	4.5 (3.0-6.2)	2.6 (1.8-3.7)	1.6 (1.1-2.2)	0.9 (0.6-1.3)	0.6 (0.4-0.8)	0.3 (0.2-0.5)	0.2 (0.1-0.3)
4	39 (31-48)	27 (20-34)	17 (13-23)	11 (7.8-15)	6.5 (4.7-9.1)	3.9 (2.8-5.5)	2.3 (1.6-3.3)	1.4 (1.0-2.0)	0.8 (0.6-1.2)	0.5 (0.4-0.7)	0.3 (0.2-0.4)
5	49 (43-57)	35 (30-43)	24 (20-30)	15 (12-20)	9.3 (7.6-13)	5.6 (4.6-7.7)	3.4 (2.7-4.6)	2.0 (1.6-2.8)	1.2 (1.0-1.7)	0.7 (0.6-1.0)	0.4 (0.4-0.6)
6	58	45	32	21	13	8.1	4.9	2.9	1.8	1.1	0.7

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Armenia.

Age = 65 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	8.3	5.3	3.4	2.1	1.3	0.8	0.5	0.3	0.2	0.1	0.1
1	12 (8.8-13)	7.6 (5.6-8.7)	4.9 (3.5-5.6)	3.1 (2.2-3.5)	1.9 (1.4-2.2)	1.2 (0.9-1.4)	0.8 (0.5-0.9)	0.5 (0.3-0.5)	0.3 (0.2-0.4)	0.2 (0.1-0.2)	0.1 (0.1-0.1)
2	16 (12-20)	11 (7.8-13)	6.9 (5.0-8.7)	4.4 (3.1-5.6)	2.8 (2.0-3.6)	1.8 (1.2-2.2)	1.1 (0.8-1.4)	0.7 (0.5-0.9)	0.4 (0.3-0.6)	0.3 (0.2-0.4)	0.2 (0.1-0.2)
3	22 (17-28)	15 (11-19)	9.7 (7.1-13)	6.3 (4.6-8.4)	4.0 (2.9-5.4)	2.5 (1.8-3.4)	1.6 (1.1-2.1)	1.0 (0.7-1.4)	0.6 (0.5-0.9)	0.4 (0.3-0.6)	0.3 (0.2-0.4)
4	29 (24-36)	20 (16-26)	13 (11-18)	8.8 (6.8-12)	5.6 (4.4-7.5)	3.6 (2.8-4.8)	2.3 (1.7-3.1)	1.4 (1.1-2.0)	0.9 (0.7-1.3)	0.6 (0.5-0.8)	0.4 (0.3-0.5)
5	36 (32-43)	26 (23-32)	18 (16-23)	12 (10-16)	7.9 (6.7-10)	5.1 (4.3-6.8)	3.2 (2.7-4.3)	2.1 (1.8-2.8)	1.3 (1.1-1.8)	0.9 (0.7-1.2)	0.6 (0.5-0.8)
6	44	33	24	16	11	7.1	4.5	2.9	1.9	1.2	0.8

Age = 70 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	6.1	4.1	2.8	1.8	1.2	0.8	0.5	0.4	0.3	0.2	0.1
1	9.4 (8.2-12)	6.4 (5.6-8.5)	4.3 (3.8-5.8)	2.9 (2.6-3.9)	1.9 (1.7-2.6)	1.3 (1.1-1.7)	0.9 (0.8-1.2)	0.6 (0.5-0.8)	0.4 (0.4-0.5)	0.3 (0.2-0.4)	0.2 (0.2-0.3)
2	14 (11-18)	9.8 (7.8-13)	6.7 (5.3-8.7)	4.6 (3.6-5.9)	3.1 (2.4-4.0)	2.1 (1.6-2.7)	1.4 (1.1-1.8)	0.9 (0.7-1.2)	0.6 (0.5-0.8)	0.4 (0.3-0.6)	0.3 (0.2-0.4)
3	20 (16-25)	15 (11-18)	10 (7.7-13)	7.0 (5.3-8.9)	4.8 (3.6-6.1)	3.2 (2.4-4.1)	2.1 (1.6-2.8)	1.5 (1.1-1.9)	1.0 (0.8-1.3)	0.7 (0.5-0.9)	0.5 (0.4-0.6)
4	28 (22-34)	21 (16-25)	15 (11-18)	11 (7.9-13)	7.3 (5.4-9.0)	5.0 (3.7-6.2)	3.3 (2.5-4.2)	2.3 (1.7-2.9)	1.6 (1.2-2.0)	1.1 (0.8-1.4)	0.7 (0.6-0.9)
5	38 (31-42)	29 (23-33)	22 (17-24)	16 (12-18)	11 (8.1-12)	7.5 (5.6-8.6)	5.1 (3.8-5.8)	3.6 (2.6-4.0)	2.5 (1.8-2.8)	1.7 (1.3-1.9)	1.2 (0.9-1.3)
6	48	39	30	22	16	11	7.8	5.4	3.8	2.7	1.8

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Armenia.

Age = 75 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	4.6	3.3	2.3	1.6	1.2	0.8	0.6	0.4	0.3	0.2	0.2
1	8.2 (5.9-17)	6.0 (4.2-13)	4.3 (3.0-9.5)	3.1 (2.2-6.8)	2.2 (1.5-4.9)	1.6 (1.1-3.5)	1.1 (0.8-2.5)	0.8 (0.6-1.8)	0.6 (0.4-1.3)	0.4 (0.3-0.9)	0.3 (0.2-0.7)
2	14 (8.2-24)	10 (5.9-19)	7.4 (4.3-14)	5.4 (3.0-10)	3.9 (2.2-7.3)	2.8 (1.5-5.2)	2.0 (1.1-3.7)	1.4 (0.8-2.7)	1.0 (0.6-2.0)	0.8 (0.4-1.4)	0.6 (0.3-1.0)
3	21 (11-31)	16 (8.3-25)	12 (6.0-19)	8.9 (4.3-14)	6.5 (3.1-10)	4.7 (2.2-7.6)	3.3 (1.6-5.5)	2.4 (1.1-4.0)	1.8 (0.8-3.0)	1.3 (0.6-2.2)	1.0 (0.4-1.6)
4	30 (15-39)	24 (11-32)	18 (8.4-25)	14 (6.1-19)	10 (4.4-14)	7.6 (3.1-10)	5.5 (2.2-7.6)	4.0 (1.6-5.7)	3.0 (1.2-4.2)	2.2 (0.9-3.1)	1.6 (0.6-2.2)
5	41 (21-47)	33 (16-39)	27 (12-32)	21 (9.0-25)	16 (6.5-19)	12 (4.7-14)	8.7 (3.4-11)	6.5 (2.5-7.9)	4.8 (1.8-5.8)	3.6 (1.3-4.3)	2.6 (1.0-3.2)
6	52	45	37	30	23	18	13	10	7.5	5.6	4.2

Age = 80 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	3.7	2.8	2.1	1.6	1.2	0.9	0.6	0.5	0.4	0.3	0.2
1	6.7 (4.7-14)	5.1 (3.6-11)	3.8 (2.7-8.6)	2.9 (2.0-6.5)	2.2 (1.5-4.9)	1.6 (1.1-3.7)	1.2 (0.8-2.8)	0.9 (0.6-2.1)	0.7 (0.5-1.6)	0.5 (0.4-1.3)	0.4 (0.3-1.0)
2	11 (6.1-20)	8.5 (4.7-16)	6.5 (3.5-12)	5.0 (2.6-9.6)	3.7 (2.0-7.3)	2.8 (1.5-5.5)	2.1 (1.1-4.2)	1.6 (0.9-3.2)	1.3 (0.7-2.5)	1.0 (0.5-1.9)	0.7 (0.4-1.5)
3	17 (8.5-27)	13 (6.5-21)	10 (4.9-17)	8.0 (3.7-13)	6.1 (2.8-10)	4.7 (2.1-7.8)	3.5 (1.6-5.9)	2.7 (1.2-4.6)	2.1 (0.9-3.6)	1.6 (0.7-2.8)	1.3 (0.6-2.1)
4	25 (12-34)	20 (8.9-28)	16 (6.8-22)	12 (5.2-18)	9.6 (3.9-14)	7.4 (2.9-11)	5.6 (2.2-8.2)	4.4 (1.7-6.5)	3.4 (1.3-5.0)	2.6 (1.0-3.9)	2.0 (0.8-3.0)
5	34 (17-40)	28 (13-33)	23 (10-27)	18 (7.7-22)	14 (5.8-17)	11 (4.4-14)	8.7 (3.4-11)	6.9 (2.6-8.3)	5.4 (2.0-6.5)	4.2 (1.6-5.1)	3.3 (1.2-3.9)
6	45	38	32	26	21	17	13	10	8.2	6.5	5.0

Ten year probability of hip fractures (%) according to BMD, the number of clinical risk factors (CRF) and age in men from Armenia.

Age = 85 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	3.0	2.4	1.8	1.4	1.1	0.9	0.7	0.6	0.4	0.4	0.3
1	5.4 (3.8-12)	4.3 (3.0-9.5)	3.4 (2.4-7.6)	2.7 (1.9-6.0)	2.1 (1.5-4.7)	1.6 (1.1-3.7)	1.3 (0.9-3.0)	1.0 (0.7-2.4)	0.8 (0.6-1.9)	0.7 (0.5-1.5)	0.5 (0.4-1.2)
2	9.1 (5.0-17)	7.3 (4.0-14)	5.8 (3.1-11)	4.6 (2.4-8.9)	3.6 (1.9-7.1)	2.9 (1.5-5.6)	2.3 (1.2-4.4)	1.8 (1.0-3.6)	1.5 (0.8-2.9)	1.2 (0.6-2.3)	1.0 (0.5-1.9)
3	14 (7.0-23)	12 (5.5-19)	9.4 (4.4-15)	7.5 (3.4-12)	6.0 (2.7-9.9)	4.7 (2.1-7.9)	3.8 (1.7-6.4)	3.1 (1.4-5.2)	2.5 (1.1-4.2)	2.0 (0.9-3.4)	1.6 (0.7-2.8)
4	21 (9.7-29)	18 (7.7-25)	14 (6.1-20)	12 (4.9-17)	9.4 (3.8-14)	7.5 (3.0-11)	6.1 (2.4-8.8)	4.9 (1.9-7.2)	4.0 (1.6-5.9)	3.3 (1.3-4.8)	2.6 (1.0-3.9)
5	30 (14-35)	25 (11-30)	21 (9.1-25)	17 (7.2-21)	14 (5.7-17)	12 (4.5-14)	9.4 (3.6-11)	7.7 (2.9-9.3)	6.3 (2.4-7.6)	5.1 (1.9-6.2)	4.2 (1.6-5.0)
6	40	35	30	25	21	17	14	12	9.6	7.9	6.5

Age = 90 years

Number of CRFs	BMD T-score (femoral neck)										
	-4.0	-3.5	-3.0	-2.5	-2.0	-1.5	-1.0	-0.5	0	0.5	1.0
0	2.0	1.6	1.3	1.1	0.9	0.7	0.6	0.5	0.4	0.4	0.3
1	3.7 (2.7-8.3)	3.1 (2.2-6.9)	2.5 (1.8-5.6)	2.1 (1.4-4.6)	1.7 (1.2-3.8)	1.4 (1.0-3.1)	1.1 (0.8-2.6)	1.0 (0.7-2.2)	0.8 (0.6-1.8)	0.7 (0.5-1.5)	0.6 (0.4-1.3)
2	6.4 (3.5-12)	5.3 (2.8-10)	4.4 (2.3-8.4)	3.6 (1.9-6.9)	2.9 (1.6-5.7)	2.4 (1.3-4.7)	2.0 (1.1-3.9)	1.7 (0.9-3.3)	1.4 (0.8-2.8)	1.2 (0.6-2.3)	1.0 (0.5-1.9)
3	10 (4.9-17)	8.6 (4.0-14)	7.2 (3.3-12)	5.9 (2.7-9.9)	4.9 (2.2-8.2)	4.0 (1.8-6.8)	3.4 (1.5-5.7)	2.9 (1.3-4.8)	2.4 (1.1-4.1)	2.0 (0.9-3.5)	1.7 (0.8-2.9)
4	16 (6.9-22)	13 (5.7-19)	11 (4.7-16)	9.4 (3.9-14)	7.8 (3.2-11)	6.5 (2.6-9.4)	5.5 (2.2-7.9)	4.6 (1.9-6.8)	3.9 (1.6-5.7)	3.3 (1.3-4.8)	2.8 (1.1-4.1)
5	23 (10-27)	20 (8.5-24)	17 (7.0-20)	14 (5.8-17)	12 (4.8-14)	10 (3.9-12)	8.6 (3.3-10)	7.3 (2.8-8.7)	6.2 (2.4-7.4)	5.3 (2.0-6.3)	4.5 (1.7-5.4)
6	32	28	25	21	18	15	13	11	9.6	8.2	7.0