

Jordan - The probabilities of a major osteoporotic fracture in women

The following tables give the 10-year probability (%) of a major osteoporotic fracture (hip, clinical spine, forearm or proximal humerus fracture) according to body mass index (BMI kg/m²), the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Jordan. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Age = 50 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 2.3 | 2.1 | 1.9 | 1.7 | 1.5 | 1.3 | 1.1 |
| 1 | 3.8 (2.6-5.7) | 3.3 (2.2-4.7) | 3.0 (2.1-4.2) | 2.6 (1.8-3.7) | 2.3 (1.6-3.2) | 2.0 (1.4-2.8) | 1.8 (1.2-2.4) |
| 2 | 6.1 (3.4-10) | 5.2 (2.8-8.7) | 4.7 (2.5-8.1) | 4.1 (2.2-7.1) | 3.6 (1.9-6.2) | 3.1 (1.6-5.4) | 2.7 (1.4-4.7) |
| 3 | 9.9 (5.0-17) | 8.2 (4.0-14) | 7.3 (3.5-13) | 6.3 (3.0-11) | 5.5 (2.6-10) | 4.8 (2.2-8.8) | 4.1 (1.9-7.7) |
| 4 | 16 (8.4-24) | 13 (7.0-20) | 11 (5.9-18) | 9.5 (5.1-16) | 8.3 (4.4-14) | 7.2 (3.8-12) | 6.2 (3.2-10) |
| 5 | 24 (14-30) | 19 (12-25) | 16 (11-22) | 14 (9.4-19) | 12 (8.2-16) | 11 (7.1-14) | 9.2 (6.2-13) |
| 6 | 36 | 28 | 24 | 21 | 18 | 16 | 14 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Jordan.

Age = 55 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 3.6 | 3.1 | 2.9 | 2.5 | 2.2 | 1.9 | 1.7 |
| 1 | 5.9 (4.2-8.6) | 5.0 (3.5-7.1) | 4.6 (3.1-6.2) | 4.0 (2.7-5.4) | 3.4 (2.3-4.7) | 3.0 (2.0-4.0) | 2.6 (1.8-3.5) |
| 2 | 9.5 (5.5-15) | 7.9 (4.4-13) | 7.0 (3.9-12) | 6.1 (3.3-10) | 5.3 (2.9-8.9) | 4.6 (2.5-7.8) | 4.0 (2.1-6.8) |
| 3 | 15 (8.3-24) | 12 (6.4-21) | 11 (5.4-19) | 9.3 (4.6-16) | 8.0 (4.0-14) | 7.0 (3.4-13) | 6.1 (3.0-11) |
| 4 | 23 (13-34) | 19 (11-28) | 16 (9.2-25) | 14 (7.9-22) | 12 (6.7-19) | 10 (5.8-17) | 9.1 (5.0-15) |
| 5 | 35 (23-42) | 28 (19-35) | 23 (16-30) | 20 (14-27) | 18 (12-23) | 15 (11-20) | 13 (9.2-18) |
| 6 | 49 | 40 | 34 | 29 | 26 | 22 | 19 |

Age = 60 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 5.1 | 4.3 | 3.9 | 3.3 | 2.9 | 2.5 | 2.2 |
| 1 | 8.2 (6.0-11) | 6.8 (4.8-9.3) | 6.0 (4.2-8.0) | 5.2 (3.6-6.9) | 4.5 (3.1-6.0) | 3.9 (2.7-5.2) | 3.4 (2.3-4.5) |
| 2 | 13 (8.0-20) | 11 (6.3-16) | 9.2 (5.3-15) | 8.0 (4.5-13) | 6.9 (3.9-11) | 6.0 (3.3-9.8) | 5.2 (2.8-8.5) |
| 3 | 20 (12-30) | 16 (9.3-26) | 14 (7.5-23) | 12 (6.4-20) | 10 (5.5-18) | 9.0 (4.7-16) | 7.8 (4.0-14) |
| 4 | 30 (19-42) | 24 (15-35) | 21 (13-31) | 18 (11-27) | 15 (9.2-24) | 13 (7.8-21) | 12 (6.7-18) |
| 5 | 43 (31-52) | 35 (25-43) | 30 (22-38) | 26 (19-33) | 22 (16-29) | 19 (14-25) | 17 (12-22) |
| 6 | 59 | 49 | 42 | 36 | 32 | 28 | 24 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Jordan.

Age = 65 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|--------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 6.8 | 5.6 | 4.9 | 4.2 | 3.6 | 3.1 | 2.7 |
| 1 | 11 (8.2-14) | 8.8 (6.5-12) | 7.6 (5.4-9.8) | 6.5 (4.6-8.4) | 5.6 (4.0-7.2) | 4.8 (3.4-6.2) | 4.2 (2.9-5.4) |
| 2 | 17 (11-24) | 14 (8.5-19) | 12 (6.9-18) | 9.9 (5.9-15) | 8.5 (5.0-13) | 7.3 (4.2-12) | 6.3 (3.6-10) |
| 3 | 25 (17-35) | 20 (13-30) | 17 (10-27) | 15 (8.5-24) | 13 (7.2-21) | 11 (6.1-18) | 9.4 (5.1-16) |
| 4 | 36 (25-48) | 30 (20-41) | 25 (17-37) | 22 (14-32) | 19 (12-28) | 16 (10-25) | 14 (8.6-21) |
| 5 | 50 (39-58) | 42 (32-50) | 36 (27-44) | 31 (23-39) | 27 (20-34) | 23 (17-30) | 20 (15-26) |
| 6 | 64 | 56 | 49 | 43 | 37 | 32 | 28 |

Age = 70 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|-------------|--------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 8.6 | 7.0 | 6.0 | 5.1 | 4.4 | 3.8 | 3.2 |
| 1 | 13 (10-16) | 11 (8.1-13) | 9.0 (6.7-11) | 7.7 (5.7-9.8) | 6.5 (4.8-8.3) | 5.5 (4.1-7.1) | 4.7 (3.5-6.1) |
| 2 | 21 (14-28) | 16 (11-22) | 13 (8.7-19) | 11 (7.3-16) | 9.7 (6.1-13) | 8.2 (5.2-11) | 7.0 (4.4-9.8) |
| 3 | 31 (22-43) | 25 (16-35) | 20 (13-30) | 17 (11-25) | 14 (9.0-22) | 12 (7.5-18) | 10 (6.3-16) |
| 4 | 45 (34-58) | 36 (27-49) | 29 (21-41) | 25 (18-35) | 21 (15-30) | 18 (12-26) | 15 (10-22) |
| 5 | 60 (54-68) | 51 (44-60) | 42 (35-51) | 36 (30-44) | 31 (25-38) | 26 (21-33) | 22 (18-28) |
| 6 | 73 | 66 | 57 | 50 | 43 | 37 | 32 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Jordan.

Age = 75 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|-------------|--------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 11 | 8.6 | 7.3 | 6.2 | 5.2 | 4.5 | 3.8 |
| 1 | 17 (12-25) | 13 (%10.0-18) | 11 (8.2-13) | 9.3 (6.9-11) | 7.8 (5.8-9.4) | 6.6 (4.9-8.1) | 5.5 (4.1-6.9) |
| 2 | 27 (17-39) | 21 (13-30) | 17 (11-23) | 14 (8.9-19) | 12 (7.5-16) | 9.8 (6.2-13) | 8.2 (5.2-11) |
| 3 | 40 (26-54) | 32 (20-46) | 25 (16-37) | 21 (13-31) | 18 (11-26) | 15 (9.2-22) | 12 (7.6-18) |
| 4 | 55 (38-68) | 47 (32-61) | 38 (26-51) | 32 (22-44) | 27 (18-37) | 22 (15-32) | 18 (12-26) |
| 5 | 68 (54-76) | 62 (48-71) | 53 (41-63) | 46 (35-55) | 39 (30-48) | 33 (25-41) | 28 (21-34) |
| 6 | 77 | 75 | 69 | 62 | 55 | 47 | 40 |

Age = 80 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|---------------|-------------|--------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 13 | 10 | 8.7 | 7.3 | 6.2 | 5.2 | 4.4 |
| 1 | 21 (15-31) | 16 (12-23) | 13 (%10.0-17) | 11 (8.3-14) | 9.2 (6.9-11) | 7.7 (5.7-9.2) | 6.5 (4.8-7.5) |
| 2 | 31 (21-45) | 25 (17-36) | 20 (13-28) | 17 (11-23) | 14 (9.0-19) | 12 (7.5-16) | 9.6 (6.2-13) |
| 3 | 44 (31-58) | 37 (25-51) | 30 (20-41) | 25 (16-34) | 21 (13-29) | 17 (11-24) | 14 (9.2-20) |
| 4 | 57 (42-69) | 51 (37-64) | 43 (30-55) | 36 (25-48) | 31 (21-41) | 26 (18-35) | 21 (15-29) |
| 5 | 69 (55-76) | 65 (51-73) | 58 (45-67) | 50 (38-59) | 43 (33-52) | 37 (27-45) | 31 (23-38) |
| 6 | 77 | 76 | 72 | 65 | 58 | 51 | 44 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Jordan.

Age = 85 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|------------|-------------|--------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 16 | 13 | 11 | 8.9 | 7.4 | 6.2 | 5.2 |
| 1 | 25 (19-37) | 20 (15-29) | 16 (12-22) | 14 (10-18) | 11 (8.4-14) | 9.3 (6.9-12) | 7.8 (5.7-9.6) |
| 2 | 37 (26-52) | 31 (21-43) | 25 (16-35) | 21 (14-29) | 17 (11-24) | 14 (9.1-19) | 12 (7.5-16) |
| 3 | 50 (35-64) | 44 (31-58) | 36 (25-49) | 31 (20-42) | 26 (17-35) | 21 (14-29) | 17 (11-24) |
| 4 | 62 (46-73) | 57 (43-69) | 50 (36-63) | 43 (31-56) | 37 (26-49) | 31 (22-42) | 26 (18-35) |
| 5 | 71 (59-78) | 69 (56-77) | 65 (51-73) | 58 (44-67) | 51 (38-60) | 44 (32-52) | 37 (27-45) |
| 6 | 78 | 78 | 76 | 71 | 65 | 58 | 50 |

Age = 90 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|------------|------------|------------|-------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 24 | 20 | 18 | 15 | 12 | 10 | 8.6 |
| 1 | 34 (26-48) | 30 (23-42) | 26 (20-35) | 22 (16-29) | 18 (14-24) | 15 (11-20) | 13 (9.2-16) |
| 2 | 45 (33-61) | 42 (30-57) | 37 (26-49) | 32 (22-42) | 27 (18-35) | 22 (15-29) | 18 (12-25) |
| 3 | 56 (41-69) | 54 (39-68) | 50 (37-62) | 43 (31-56) | 37 (26-49) | 32 (21-42) | 26 (18-35) |
| 4 | 65 (51-76) | 64 (50-75) | 62 (49-72) | 56 (43-67) | 49 (36-61) | 43 (31-54) | 37 (26-48) |
| 5 | 73 (62-79) | 73 (62-79) | 72 (61-78) | 67 (55-74) | 61 (49-69) | 55 (42-63) | 48 (36-57) |
| 6 | 79 | 79 | 79 | 75 | 71 | 65 | 59 |