



Lebanon - The probabilities of a major osteoporotic fracture in women

The following tables give the 10-year probability (%) of a major osteoporotic fracture (hip, clinical spine, forearm or proximal humerus fracture) according to body mass index (BMI kg/m²), the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Lebanon. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Age = 50 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 1.1 | 1.0 | 1.0 | 0.8 | 0.7 | 0.6 | 0.6 |
| 1 | 1.9 (1.3-2.9) | 1.6 (1.1-2.4) | 1.5 (1.0-2.1) | 1.3 (0.9-1.8) | 1.1 (0.8-1.6) | 1.0 (0.7-1.4) | 0.9 (0.6-1.2) |
| 2 | 3.1 (1.7-5.2) | 2.6 (1.4-4.4) | 2.4 (1.3-4.1) | 2.1 (1.1-3.6) | 1.8 (0.9-3.1) | 1.6 (0.8-2.7) | 1.4 (0.7-2.4) |
| 3 | 5.1 (2.5-8.7) | 4.2 (2.0-7.4) | 3.7 (1.7-6.7) | 3.2 (1.5-5.9) | 2.8 (1.3-5.1) | 2.4 (1.1-4.5) | 2.1 (1.0-3.9) |
| 4 | 8.1 (4.3-13) | 6.5 (3.5-10) | 5.6 (3.0-9.3) | 4.8 (2.5-8.0) | 4.2 (2.2-7.0) | 3.6 (1.9-6.1) | 3.1 (1.6-5.3) |
| 5 | 13 (7.5-16) | 10 (6.2-13) | 8.5 (5.5-11) | 7.3 (4.8-9.8) | 6.3 (4.1-8.5) | 5.4 (3.6-7.4) | 4.7 (3.1-6.4) |
| 6 | 20 | 15 | 13 | 11 | 9.4 | 8.1 | 7.0 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Lebanon.

Age = 55 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 1.9 | 1.6 | 1.5 | 1.3 | 1.1 | 1.0 | 0.9 |
| 1 | 3.1 (2.2-4.6) | 2.6 (1.8-3.7) | 2.4 (1.6-3.3) | 2.1 (1.4-2.8) | 1.8 (1.2-2.4) | 1.6 (1.1-2.1) | 1.3 (0.9-1.8) |
| 2 | 5.1 (2.9-8.1) | 4.2 (2.3-6.7) | 3.7 (2.0-6.2) | 3.2 (1.7-5.4) | 2.8 (1.5-4.7) | 2.4 (1.3-4.1) | 2.1 (1.1-3.6) |
| 3 | 8.1 (4.4-13) | 6.6 (3.4-11) | 5.7 (2.8-10) | 4.9 (2.4-8.8) | 4.2 (2.1-7.7) | 3.7 (1.8-6.7) | 3.2 (1.5-5.8) |
| 4 | 13 (7.2-19) | 10 (5.9-16) | 8.7 (4.9-14) | 7.5 (4.2-12) | 6.4 (3.5-10) | 5.5 (3.0-9.1) | 4.8 (2.6-7.9) |
| 5 | 20 (13-25) | 16 (10-20) | 13 (8.8-17) | 11 (7.6-15) | 9.6 (6.5-13) | 8.3 (5.6-11) | 7.1 (4.9-9.6) |
| 6 | 30 | 23 | 19 | 16 | 14 | 12 | 10 |

Age = 60 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 3.2 | 2.7 | 2.4 | 2.1 | 1.8 | 1.6 | 1.4 |
| 1 | 5.1 (3.7-7.2) | 4.3 (3.0-5.8) | 3.8 (2.6-5.1) | 3.3 (2.3-4.4) | 2.8 (1.9-3.8) | 2.4 (1.7-3.2) | 2.1 (1.5-2.8) |
| 2 | 8.2 (5.0-13) | 6.7 (3.9-10) | 5.9 (3.3-9.4) | 5.0 (2.8-8.2) | 4.3 (2.4-7.1) | 3.8 (2.1-6.2) | 3.2 (1.8-5.4) |
| 3 | 13 (7.7-20) | 10 (5.8-17) | 8.9 (4.7-15) | 7.7 (4.0-13) | 6.6 (3.4-12) | 5.7 (2.9-10) | 4.9 (2.5-8.7) |
| 4 | 20 (12-28) | 16 (9.8-24) | 13 (8.1-21) | 11 (6.9-18) | 9.9 (5.8-16) | 8.5 (4.9-14) | 7.3 (4.2-12) |
| 5 | 30 (20-36) | 24 (16-30) | 20 (14-25) | 17 (12-22) | 15 (10-19) | 13 (8.9-17) | 11 (7.7-14) |
| 6 | 43 | 34 | 28 | 25 | 21 | 18 | 16 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Lebanon.

Age = 65 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|---------------|---------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 5.0 | 4.2 | 3.7 | 3.2 | 2.7 | 2.3 | 2.0 |
| 1 | 8.0 (6.0-11) | 6.5 (4.8-8.6) | 5.7 (4.1-7.4) | 4.9 (3.5-6.3) | 4.2 (2.9-5.4) | 3.6 (2.5-4.7) | 3.1 (2.2-4.0) |
| 2 | 12 (8.1-18) | 10 (6.3-15) | 8.7 (5.2-13) | 7.5 (4.4-12) | 6.4 (3.7-10) | 5.5 (3.1-8.8) | 4.7 (2.7-7.6) |
| 3 | 19 (13-27) | 15 (9.5-23) | 13 (7.5-21) | 11 (6.3-18) | 9.6 (5.3-16) | 8.2 (4.5-14) | 7.1 (3.8-12) |
| 4 | 28 (19-38) | 23 (15-32) | 19 (13-29) | 17 (11-25) | 14 (9.0-22) | 12 (7.5-19) | 10 (6.4-16) |
| 5 | 40 (30-47) | 33 (25-40) | 28 (21-35) | 24 (18-30) | 21 (15-26) | 18 (13-23) | 15 (11-20) |
| 6 | 53 | 45 | 39 | 34 | 29 | 25 | 22 |

Age = 70 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|-------------|--------------|---------------|---------------|---------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 8.2 | 6.8 | 6.0 | 5.1 | 4.4 | 3.7 | 3.2 |
| 1 | 13 (9.6-16) | 10 (7.8-13) | 9.0 (6.7-11) | 7.6 (5.6-9.7) | 6.4 (4.7-8.3) | 5.5 (4.0-7.1) | 4.7 (3.4-6.1) |
| 2 | 20 (13-27) | 16 (10-22) | 13 (8.6-18) | 11 (7.2-16) | 9.5 (6.0-13) | 8.1 (5.1-11) | 6.8 (4.3-9.6) |
| 3 | 29 (20-40) | 24 (16-34) | 20 (13-29) | 17 (11-25) | 14 (8.8-21) | 12 (7.4-18) | 10 (6.2-15) |
| 4 | 42 (31-54) | 35 (25-47) | 29 (21-40) | 24 (17-35) | 21 (14-30) | 17 (12-25) | 15 (10-21) |
| 5 | 55 (49-64) | 48 (42-57) | 41 (34-49) | 35 (29-43) | 30 (24-37) | 25 (20-32) | 21 (17-27) |
| 6 | 67 | 63 | 55 | 48 | 42 | 36 | 30 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Lebanon.

Age = 75 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|-------------|-------------|--------------|---------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 13 | 11 | 10 | 8.5 | 7.2 | 6.1 | 5.2 |
| 1 | 21 (15-30) | 18 (13-23) | 15 (11-18) | 13 (9.3-15) | 11 (7.8-13) | 8.9 (6.5-11) | 7.5 (5.5-9.3) |
| 2 | 31 (20-45) | 27 (17-38) | 22 (14-30) | 19 (12-25) | 16 (10-21) | 13 (8.4-18) | 11 (7.0-15) |
| 3 | 43 (28-59) | 38 (25-52) | 33 (21-45) | 28 (18-39) | 23 (15-33) | 19 (12-27) | 16 (10-23) |
| 4 | 56 (40-68) | 52 (36-65) | 46 (32-59) | 39 (27-52) | 33 (22-45) | 28 (19-39) | 23 (16-33) |
| 5 | 66 (54-74) | 64 (52-73) | 60 (48-69) | 53 (42-63) | 46 (36-56) | 40 (31-49) | 34 (26-42) |
| 6 | 75 | 74 | 72 | 67 | 60 | 53 | 46 |

Age = 80 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|------------|------------|-------------|-------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 19 | 17 | 15 | 13 | 11 | 9.1 | 7.6 |
| 1 | 28 (20-40) | 25 (19-35) | 22 (17-29) | 19 (14-24) | 16 (12-20) | 13 (9.5-16) | 11 (7.9-13) |
| 2 | 38 (26-53) | 35 (25-48) | 32 (22-42) | 27 (18-35) | 23 (15-30) | 19 (12-25) | 16 (10-21) |
| 3 | 49 (34-63) | 47 (33-60) | 43 (31-56) | 37 (26-49) | 32 (21-42) | 27 (18-36) | 22 (15-30) |
| 4 | 59 (44-70) | 58 (44-69) | 55 (42-67) | 49 (36-61) | 43 (30-54) | 36 (25-47) | 31 (21-41) |
| 5 | 68 (56-75) | 67 (55-74) | 66 (55-74) | 60 (49-69) | 54 (42-63) | 48 (37-57) | 41 (31-50) |
| 6 | 76 | 75 | 75 | 70 | 65 | 59 | 52 |

Ten year probability of osteoporotic fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in women from Lebanon.

Age = 85 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|------------|------------|------------|------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 23 | 22 | 21 | 17 | 14 | 12 | 9.8 |
| 1 | 33 (24-46) | 31 (23-44) | 30 (22-40) | 25 (18-34) | 21 (15-28) | 17 (12-23) | 14 (10-19) |
| 2 | 43 (30-58) | 42 (30-56) | 40 (29-53) | 34 (24-46) | 29 (20-40) | 24 (16-33) | 20 (13-28) |
| 3 | 53 (39-67) | 52 (37-66) | 51 (37-65) | 45 (31-58) | 39 (26-52) | 33 (21-45) | 28 (18-39) |
| 4 | 63 (49-74) | 62 (48-73) | 61 (48-72) | 55 (42-67) | 49 (36-61) | 43 (30-54) | 37 (25-48) |
| 5 | 71 (60-76) | 71 (59-77) | 70 (59-76) | 65 (53-72) | 59 (47-67) | 53 (41-62) | 47 (36-56) |
| 6 | 75 | 77 | 77 | 73 | 68 | 63 | 57 |

Age = 90 years

| Number of CRFs | BMI (kg/m ²) | | | | | | |
|----------------|--------------------------|------------|------------|------------|------------|------------|------------|
| | 15 | 20 | 25 | 30 | 35 | 40 | 45 |
| 0 | 27 | 26 | 25 | 20 | 17 | 14 | 11 |
| 1 | 37 (29-52) | 36 (27-49) | 35 (26-47) | 29 (21-40) | 24 (17-34) | 20 (14-28) | 16 (11-23) |
| 2 | 48 (35-63) | 46 (34-61) | 45 (33-59) | 39 (27-53) | 33 (22-46) | 28 (18-40) | 23 (15-33) |
| 3 | 58 (44-70) | 57 (42-70) | 55 (41-69) | 49 (35-63) | 43 (29-57) | 37 (24-51) | 32 (20-44) |
| 4 | 66 (54-75) | 66 (53-75) | 65 (52-75) | 59 (46-70) | 53 (39-65) | 47 (34-59) | 41 (28-53) |
| 5 | 69 (63-73) | 72 (63-76) | 72 (63-78) | 68 (57-75) | 63 (51-71) | 57 (45-66) | 51 (39-60) |
| 6 | 67 | 74 | 77 | 75 | 71 | 67 | 61 |